

NCAP EIP 2023 audit: Analysis plan

Analysis to be conducted on all scoring matrix items at a national, Trust, regional and team level. Results will be displayed via an online dashboard.

	Scoring matrix item	Question numbers in casenote audit tool	Analysis plan
1	Percentage of service users with first episode psychosis that were allocated to, and engaged with, an EIP care coordinator within 2 weeks of receipt of referral	1.5, 1.6 & 1.7	Numerator: All those who were allocated to and engaged with an EIP care coordinator within two weeks of referral Denominator : All
2.1	Percentage of service users with first episode psychosis that took up Cognitive Behavioural Therapy for psychosis (CBTp)	2.1	Numerator: All those who commenced a course of CBTp, delivered by a person with relevant skills, experience and competencies (answered 'took up' CBTp) Denominator: all
2.2	Percentage of service users with first episode psychosis that took up supported employment and education programmes	2.2.1 & 2.2.2	 Numerator: all those who have commenced a supported employment programme (Answered 'took up'). Denominator: all those who are not in work, education, or training at the time of their initial assessment (answered 'No')
2.3	Percentage of service users with first episode psychosis and their families that took up Family Interventions	2.3 & 2.4.1	Numerator: All those who commenced a course of FI, delivered by a person with relevant skills, experience, and competencies (answered 'took up' FI)



	Scoring matrix item	Question numbers in casenote audit tool	Analysis plan
			Denominator : all those who have an identified family member, friend or carer who supports them (answered 'yes' and are happy to be contacted)
2.4	Percentage of carers that took up carer-focussed education and support programmes	2.4.1 & 2.4.2	 Numerator: all carers who have commenced a course of a carer-focused education and support programme (answered 'yes'). Denominator: carers from all those who have an identified family member, friend or carer who supports them (answered 'yes' and those who selected that they did not want this person to be contacted)
2.5	Percentage of service users with first episode psychosis that have had a physical health review and relevant interventions in the last year Includes the following physical health reviews: • Smoking status • Alcohol intake • Substance misuse • BMI • Blood pressure • Glucose • Cholesterol	2.5.1-2.5.14	For each individual physical health review: Numerator: all those who received a screening (including refusals/those who were pregnant or gave birth in last 6 weeks) Denominator: all For all 7 physical health reviews collectively: Numerator : all those who received all seven individual PH measures (including refusals/those who were pregnant or gave birth in last 6 weeks) Denominator : all For each individual physical health intervention: Numerator: all those offered a relevant intervention (including refusals) Denominator : all who required an intervention as per the Lester tool



	Scoring matrix item	Question numbers in casenote audit tool	Analysis plan
	 And following psychical health interventions: Smoking cessation Harmful alcohol use Substance misuse Weight gain/obesity Hypertension Diabetes/high risk of diabetes Dyslipidaemia 		To meet this standard service users must have been offered all 7 physical health screenings and offered a relevant intervention where screening indicated as determined by the Lester tool. Composite measure of the above to determine the overall item score: Numerator: all those who received screenings (including refusals/those who were pregnant or gave birth in last 6 weeks) and any relevant interventions for all seven PH measures (including refusals) Denominator: all
3	Percentage of service users for whom two or more outcome measures (from HoNOS/HoNOSCA, DIALOG and QPR) were recorded at least twice (assessment and one other time point)	3.1-3.5	Numerator: all those with 2 or more outcome measures recorded more than once (DIALOG and any other outcome measure) Denominator: all



	Additional Analysis	Relevant question numbers	Analysis plan
A.1	Percentage of service users with first episode psychosis who have not responded adequately to or tolerated treatment with at least two antipsychotic drugs are offered clozapine	4.1 & 4.2	 Numerator: all those who have been offered clozapine (accepted or refused) Denominator: all those who have not responded adequately to or tolerated treatment with at least 2 antipsychotic drugs