

NCAP EIP 2019/2020 audit standards and outcome indicator (Wales)



Standards	
S1	Service users with first episode of psychosis start treatment in early intervention in psychosis services within two weeks of referral (allocated to, and engaged with, an EIP care coordinator).
S2	Service users with first episode psychosis take up Cognitive Behavioural Therapy for psychosis (CBTp).
S3	Service users with first episode psychosis and their families take up Family Interventions.
S4	Service users with first episode psychosis who have not responded adequately to or tolerated treatment with at least two antipsychotic drugs are offered clozapine.
S5	Service users with first episode psychosis take up supported employment and education programmes.
S6	Service users receive a physical health review annually. This includes the following measures: <ul style="list-style-type: none"> • Smoking status • Alcohol intake • Substance misuse • BMI • Blood pressure • Glucose • Cholesterol
S7	Service users are offered relevant interventions for their physical health for the following measures: <ul style="list-style-type: none"> • Smoking cessation • Harmful alcohol use • Substance misuse • Weight gain/obesity • Hypertension • Diabetes/high risk of diabetes • Dyslipidaemia
S8	Carers take up or are referred to carer-focused education and support programmes.

Outcome indicator	
I.1	Clinical outcome measurement data for service users (two or more outcome measures from DIALOG, Other) is recorded at least twice (assessment and one other time point).