


## NCAP EIP audit 2020/2021 standards and outcome indicator



Standards	
<b>S1*</b>	Service users with first episode of psychosis start treatment in early intervention in psychosis services within two weeks of referral (allocated to, and engaged with, an EIP care coordinator).
<b>S2</b>	Service users with first episode psychosis take up Cognitive Behavioural Therapy for psychosis (CBTp).
<b>S3</b>	Service users with first episode psychosis and their families take up Family Interventions.
<b>S4</b>	Service users with first episode psychosis who have not responded adequately to or tolerated treatment with at least two antipsychotic drugs are offered clozapine.
<b>S5</b>	Service users with first episode psychosis take up supported employment and education programmes.
<b>S6</b>	Service users receive a physical health review annually. This includes the following measures: <ul style="list-style-type: none"> <li>• Smoking status</li> <li>• Alcohol intake</li> <li>• Substance misuse</li> <li>• BMI</li> <li>• Blood pressure</li> <li>• Glucose</li> <li>• Cholesterol</li> </ul>
<b>S7</b>	Service users are offered relevant interventions for their physical health for the following measures: <ul style="list-style-type: none"> <li>• Smoking cessation</li> <li>• Harmful alcohol use</li> <li>• Substance misuse</li> <li>• Weight gain/obesity</li> <li>• Hypertension</li> <li>• Diabetes/high risk of diabetes</li> <li>• Dyslipidaemia</li> </ul>
<b>S8</b>	Carers take up carer-focused education and support programmes.

Outcome indicator	
<b>I.1</b>	Clinical outcome measurement data for service users (two or more outcome measures from HoNOS/HoNOSCA, DIALOG, QPR) is recorded at least twice (assessment and one other time point).

\*data will not be collected for this standard, the Early Intervention in Psychosis Waiting Times data published by NHS Digital will be used.