


Managing Alcohol Related Brain Damage in the Community

A developing service

Cheshire and Wirral Partnership NHS Foundation Trust
Mersey Care NHS Trust
University of Liverpool

ARBD



Prevalence

post mortem studies (RCP report)

- ▶ 30% of alcohol misusers have ARD or cerebella damage (thiamine deficient)
- ▶ Diagnostic problems
 - 18% of patients diagnosed during their life time
 - 10% of WK patients have classical signs
- ▶ Estimated under diagnosis of 90%
- ▶ ~1.5% of patients of the general population at post mortem

summary findings

Boughy et al (2007), MacRae R, Cox S. (2003)

- ▶ ARBD is stigmatised;
 - Self inflicted
 - Low priority
 - Double jeopardy; dementia and alcohol dependency
- ▶ Lack of professional responsibility for care
- ▶ Lack of training in medical professions
- ▶ Evidence of no clear clinical pathway
- ▶ Lack of commissioning of care

The case for rehabilitation

Evidence drawn from brain injury, ARBD and alcohol treatment studies (no RCTs)

- ▶ There is likely to be variable cognitive recovery associated with abstinence provided the environment is correct.
- ▶ Head injury and ARBD studies indicate that:
 - Specific cognitive training enhances recovery and is generalisable to other cognitive domains.
 - Ecologically driven rehabilitation is beneficial.
- ▶ Patients receiving 'specialised services' fare better than those receiving generic services

Identifying cases

Screening for referral to the ARBD service
from acute hospital wards

All three criteria should be positive

1. Probable history of heavy, long standing alcohol drinking: 35 units or more a week for at least five years.
2. Confusion, memory problems, doubt about capacity and concerns about risk on discharge, after withdrawal/physical stabilisation.
3. Three or more admissions into hospital and/or A&E in one year probably associated either directly (withdrawal, unconscious) or indirectly (trauma, organ disease, etc.) with alcohol ingestion.

Or

One or more delayed discharges from general hospital wards in the last 12 months. (a delayed discharge is defined as patients staying on the acute medical/surgical ward because of social and/or psychiatric problems).

(Adapted through pilot work from Oslin's criteria relating to diagnosis of alcohol related dementia. (Oslin DW, Carey MS. Alcohol related dementia; Validation of diagnostic criteria. American Journal of Geriatric Psychiatry 2003;11(4):441-7.))

Manual development

- ▶ Funded by an innovation grant (NW SHA)
- ▶ Informed by evidence relating to rehabilitation of people with
 - Acquired brain injury
 - Alcohol related brain damage
 - Alcohol misuse and dependency
- ▶ Designed for non specialised community teams working with non specialised nursing homes, community institutions and support services.
- ▶ Developed through:
 - Small group education
 - Supervision of cases
 - Feedback through in depth interviews with patients

Case study 1

- ▶ 42 year old women
- ▶ Found by police , unconscious, in a skip after sexual assaults
- ▶ No relatives
- ▶ Long standing drinking
- ▶ Multiple hospital admissions in acute medical/surgical hospitals.
- ▶ Protracted malnutrition and chronic liver disease
- ▶ Previous periods of WE a year ago
- ▶ Admitted to medical ward
 - GGT >1000
- ▶ Physically stabilised: Psychiatry liaison; referred to service.

Case study 1

Neuropsychological profile

Quite incapacitated with no insight

- ▶ Profound problems in learning new information
- ▶ Retrograde amnesia
- ▶ Varying confabulation
- ▶ Processing speed much reduced: leading to missing information
- ▶ Perceptual organisation low; 'difficulty in seeing the whole picture'
- ▶ Graded naming was good
- ▶ Problem solving tests (behavioural assessment of dysexecutive syndrome): proneness to break rules, problems in planning
- ▶ Sentence completion is very poor due to inability to inhibit responses.

Case study 1

Progress after hospital

- ▶ Phase 2: CTO to specialised rehabilitation unit 6 months (out of area).
- ▶ Phase 2/3:
 - Case specific commissioned package of care
 - Placed in (Non specialised) mental health sheltered accommodation with additional support supervised rehab through team (CTO).
 - Very good progress, cognitive impairments improved, self catering in structured environment. Alcohol free
- ▶ General improvement in independence and memory.
Rehabilitative support withdrawn
- ▶ Quality of life good, CTO closed
- ▶ Still requires supervised care but now able to shop and manage on a day to day basis

Case study 2

- ▶ 55 year old patient transferred from Liverpool to a house rented by his brother in Wirral so the patient can be looked after by his brother.
- ▶ Admitted to acute medical care as a consequence of uncontrolled convulsions.
- ▶ In an acute confusional state, went through withdrawal, GGT: 820
- ▶ ACE-R score 23 (out of 100)
- ▶ Transferred from acute hospital to supported living for people with mental illness with additional carer support.

Case study 2

- ▶ Long term drinking with Many years of cognitive dysfunction.
- ▶ No capacity or awareness of effects of alcohol on convulsions/cognitive state or physical state
- ▶ Highly suggestible with little reasoning and proneness to confabulate
- ▶ No cognitive capacity to manage personal funding
- ▶ Significant esophageal varicies and peripheral neuropathy
- ▶ Has little or no memories of previous few hours or recent events.
- ▶ Pre-occupation with alcohol access

Case study 2

Two years later:

- ▶ ACE-R score 56; (23)
- ▶ Convulsions stabilised
- ▶ Settled in supported living
- ▶ Drinking two cans of lager a day
- ▶ Going out with brother regularly
- ▶ One hospital admission over night for convulsion

Current status of rehab.

- ▶ Activity scheduling
- ▶ Targeted graded task assignment

Problems

Still no capacity to understand alcohol related problems:

Cannot retain information

Cannot understand what effects alcohol has on his physical and cognitive state

Is very suggestible and vulnerable

Finances still controlled by Brother, still engaged with advocate.

Service design and delivery

Objectives:

- To improve quality of life of ARBD sufferers
- To reduce admissions into acute medical/surgical care
 - To develop local skills and services

Clinical review of consecutive series of patients referred

(up to end of 2010)

- ▶ Population size: 300,000
- ▶ The additional members of the team:
 - One band six nurse
 - One band 6 social worker (AMP)
- ▶ Approximately 2 patients a month
- ▶ 41 cases accepted and followed up for an average of 25 months (1–131)

Demographic and clinical information

- ▶ 11 female (average age 50.1)
- ▶ 30 males (average age: 59.9)
- ▶ All had experienced acute confusional states: Encephalopathy, delirium tremens/withdrawal confusion.
- ▶ 4 were admitted through medical intensive care; warranting intubation

Medical and psychiatric presentation/history

History of, or presentation with co-morbid physical conditions	Number of patients	History of, or presenting co-morbid mental illness	Number of patients
Unspecified encephalopathy	8	Depression	17
Convulsions	10	Aggression	8
Peripheral neuropathy	8	Cerebral ischaemic/infarcts	9
Upper motor neurone signs	3	Subdurals/ significant head trauma or anoxic brain damage	6
Cerebella signs	4	polydypsia	1
History of portal hypertension/oesophageal varicies	4	Bipolar affective disorder	1
Deep venous thrombosis	4	PTSD	1
Diabetes	4	Hoarding	1
Chronic urinary incontinence/renal disease	4	Heroine dependency (on methadone)	1
Hepatitis C positive	2		
Heart failure/fibrillation	5		
Pancreatic disease	2		
Duodenitis/gastritis/ulcers	5		
History of significant fractures/dislocations	6		

- Cerebro-vascular disease/head trauma in 1/3cases
- 1 patient went through PICU
- 5 patients on CTOs and one on guardianship

Cognitive profile

- ▶ ACE-R was carried out on a consecutive series of 22 of the 41 patients
 - Conducted at of an average of 8 months into rehab programme: (3–12 months)
 - group average ACE-R score was 65.7 (range: 30–93, maximum score; 100)
 - Group average Mini Mental State Examination score (incorporated within the ACE-R) was 23.3 (range 11–30; maximum score 30)
 - the individual with a score of 30 on the Mini Mental State Examination had a score of 93 on the ACE-R. This individual had clearly defined difficulty in higher order reasoning and memory. The patient is currently held under guardianship of the mental health act.

HoNOS-ABI cognitive profile (n=38)

▶ First three months (n=26)

- 10 : severe cognitive problems; hardly capable of the simplest tasks, with very poor memory, attentional problems and hardly able to learn new information.
- 13: suffering from moderate cognitive problems; including disorientation, attentional problems, organisational difficulties and difficulty in thinking clearly.
- 3 had either circumscribed cognitive deficits or milder impairment.

▶ 3 months to one year (n=12)

- 2: severe cognitive impairment.
- 4: patients had moderate cognitive impairment
- 6: milder cognitive impairment characterised by disorientation, difficulty in prioritising tasks, organisational issues and definite problems in learning new information.
- 5: severe or moderate degrees of hallucinations, delusions or confabulation and another.
- 5: milder, less intrusive symptoms of a similar nature.

Clinical follow-up and outcomes

16 assessment

11 in institutional care
All abstinent
(1 of which died)

5 supported at Home of which
2 abstinent
1 controlled drinking
2 relapsed: 1 in a com. withdrawal facility

8 rehabilitation

6 in institutional care
All abstinent

1 rehabilitated at Home: abstinent

1 lost to follow-up
Assumed uncontrolled drinking

17: adaptation/relapse prevention

5 in nursing homes

3 remain in NH
Co-Morbid MI

2 at home
both died
(abstinent)

8 abstinent
1 died

2 controlled drinking

2 relapsed;
uncontrolled drinking

HoNOS-ABI outcomes of 12 of the 17 patients in last phases of rehab

- ▶ Group average rating scores fell in domains active disturbance of social behaviour
 - problem drinking and drug use
 - cognitive problems
 - physical illness and disability
 - experience of hallucinations
 - delusions and confabulation
 - problems with relationships
 - problems with activities of daily living
 - problems with living conditions and problems with activities
 - No patients were rated as experiencing self directed injury
- ▶ There was an increase in group ratings of the experience of depression and other mental conditions.

Unplanned hospital admissions

- ▶ 5 years preceding end of index admission:
 - 205 patient years
 - 41 patients had 4418 days of admission
 - 0.53 acute medical/surgical bed days per patient each patient year
- ▶ 41 patients were followed up for 85.6 patient-years
 - 295 days of inpatient care in acute medical or surgical wards
 - 0.08 acute medical/surgical inpatient days per patient each patient-year
- ▶ Reduction of acute medical surgical beds by 85%

Summary outcomes

- ▶ N=41
- ▶ 32 non relapsed, surviving patients:
 - 18 reside in community institutions
 - 14 are supported in their own home
- ▶ Of the remaining 9 patients
 - 10% mortality rate
 - 5 in uncontrolled drinking

General points

- ▶ Rehabilitation best carried out by Neuro-rehabilitation but:
 - It can be carried out by teams that are familiar with assessment of cognitive damage
 - It can be carried out by non trained staff under supervision
- ▶ Cost of service delivery can in part be offset against reduction in acute medical inpatient stay
- ▶ Commissioning of individual care packages can play an important role
- ▶ Care packages need on-going review (and usually cost less as the rehabilitation progresses)
- ▶ Frequent use of Mental Incapacity act and Mental Health act.
- ▶ Development of 'in-reach' required
- ▶ Relatively active follow-up required
- ▶ 'Specialist' nursing homes/insititutions not required.
 - Use market forces to
 - build expertise
 - develop patient specific care plans

For further information (copy of manual, brief lit. review etc)
Please contact Prof Ken Wilson: k.c.m.wilson@liverpool.ac.uk