

NCAP EIP 2021-22 audit standards and outcome indicator - Ireland

	Standards Standa		
S1	Service users with first episode of psychosis start treatment in early intervention in psychosis services within two		
	weeks of referral (allocated to, and engaged with, an EIP care coordinator).		
S2	Service users with first episode psychosis take up Cognitive Behavioural Therapy for psychosis (CBTp).		
S3	Service users with first episode psychosis and their families take up Family Interventions.		
S4	Service users with first episode psychosis who have not responded adequately to or tolerated treatment with at least		
	two antipsychotic drugs are offered clozapine.		
S5	Service users with first episode psychosis take up supported employment and education programmes.		
S6	Service users receive a physical health review annually. This includes the following measures:		
	Smoking status		
	Alcohol intake		
	Substance misuse		
	• BMI		
	Blood pressure		
	Glucose		
	• Cholesterol		
S7	Service users are offered relevant interventions for their physical health for the following measures:		
	Smoking cessation		
	Harmful alcohol use		
	Substance misuse		
	Weight gain/obesity		
	Hypertension		
	Diabetes/high risk of diabetes		
	Dyslipidaemia		
S8	Carers take up carer-focused education and support programmes.		

		Outcome indicator
I	.1	Clinical outcome measurement data for service users (two or more outcome measures) is recorded at least twice
		(assessment and one other time point).